arXiv:hep-ph/0606092v2 18 Jul 2006

Role of electromagnetic dipole operator in the electroweak penguin dominated vector meson decays of B meson*

Cai-Dian Lü[a,b],Yue-Long Shen[b] and Wei Wang[b]
a CCAST (World Laboratory), P.O. Box 8730, Beijing 100080, P.R. China
b Institute of High Energy Physics, CAS, P.O.Box 918(4), 100049, P.R. China†

The pure annihilation type decays $B_d^0 \to \phi\gamma$ and $B_s \to \rho\gamma$ receive only color suppressed penguin contributions with a very small branching ratio in the standard model. When we include the previously neglected electromagnetic dipole operator, the branching ratios can be enhanced one order magnitude larger than previous study using QCD factorization approach. That is $\mathcal{BR}(\bar{B}_d^0 \to \phi\gamma) \simeq 1 \times 10^{-11}$ and $\mathcal{BR}(B_s \to \rho\gamma) \sim (6-16) \times 10^{-10}$. The new effect can also give a large contribution, of order 10^{-9}, to transverse polarization of $B \to \phi\rho$ and $B \to \omega\phi$ which is comparable to the longitudinal part. These effects can be detected in the LHCb experiment and the Super-B factories.

PACS numbers:

The charmless two-body B decays provide rigorous information on flavor changing phenomena in the weak interactions of quarks, so it has attracted much attention to check the consistency of the standard model (SM) and to explore the existence of the possible new physics beyond standard model. On the other hand, the rare B decays also serve as a laboratory for hadronic dynamics. However, predictions for many interesting decays are always polluted by the hadronization, which have hindered us in extracting weak interaction information precisely from the available measurements. To handle the decay processes, many factorization approaches have been developed to separate nonperturbative dynamics from the perturbative part, such as naive factorization [1], generalized factorization [2, 3], the QCD factorization (QCDF) [4], perturbative QCD approach (PQCD) [5] and the soft-collinear effective theory (SCET) [6].

Two vector decays of B meson shed light on the helicity structure of weak interactions through polarization studies. The naive counting rules [7] based on the factorization approaches predict that the longitudinal polarization dominates the decay ratios and the transverse polarizations are suppressed due to the helicity flips of the quark in the final state hadrons. But for penguin-dominated decays such as $B \to \phi K^*$ and $B \to \rho K^*$ [8], large transverse polarizations are observed.

---

† Mailing address

*This work is partly supported by National Science Foundation of China under contract No. 10475085



FIG. 1: New contributions to $B \to VV$ decays

Beneke, Rohrer and Yang [9] discussed a novel electromagnetic penguin contribution to the transverse helicity amplitudes, which is shown in Fig.1. The neutral vector meson generated by the photon is mainly transversely polarized. Though suppressed by the fine structure constant α_{em}, it is enhanced over the leading terms by m_B/Λ. So the transverse helicity amplitude of $B \to \rho^0 K^*$ can be enhanced by this effect. The validity of this effect can be tested in the ratio of transverse polarization for $B \to \rho^0 K^*$ over $B \to \rho^+ K^*$ in the near future.

It may play an important role in the electroweak penguin dominated vector meson decays of B meson. In the standard model, it is suppressed to produce a ϕ meson through strong interaction for three or more gluons are needed which are called hairpin diagrams [10] as shown in Fig. 2. So the electromagnetic dipole operator may become important in $B \to \phi\gamma$ and $B \to \phi\rho$ decays. We examine the new effect in these modes below.

In the standard model, the effective weak Hamiltonian mediating flavor-changing neutral current transitions of the type $b \to D (D = d, s)$ has the form [11]:

$$\mathcal{H}_{eff} = \frac{G_F}{\sqrt{2}}\Big[\sum_{p=u,c} V_{pb}V^*_{pD}\Big(C_1 O_1^p + C_2 O_2^p\Big) - V_{tb}V^*_{tD}\sum_{i=3}^{10}\Big(C_i O_i + C_{7\gamma}O_{7\gamma} + C_{8g}O_{8g}\Big)\Big]. \tag{1}$$

For convenience, we specify the below operators in $\mathcal{H}_{eff}$:

$$\begin{aligned}
O_1^p &= \bar{p}_\beta\gamma^\mu L b_\alpha \cdot \bar{D}_\alpha\gamma_\mu L p_\beta, \quad O_2^p = \bar{p}_\alpha\gamma^\mu L b_\alpha \cdot \bar{D}_\beta\gamma_\mu L p_\beta, \\
O_3 &= \bar{D}_\alpha\gamma^\mu L b_\alpha \cdot \sum_q \bar{q}_\beta\gamma_\mu L q_\beta, \quad O_4 = \bar{D}_\beta\gamma^\mu L b_\alpha \cdot \sum_q \bar{q}_\alpha\gamma_\mu L q_\beta, \\
O_5 &= \bar{D}_\alpha\gamma^\mu L b_\alpha \cdot \sum_q \bar{q}_\beta\gamma_\mu R p_\beta, \quad O_6 = \bar{D}_\beta\gamma^\mu L b_\alpha \cdot \sum_q \bar{q}_\alpha\gamma_\mu R q_\beta, \\
O_7 &= \frac{3}{2}\bar{D}_\alpha\gamma^\mu L b_\alpha \cdot \sum_q e_q\bar{q}_\alpha\gamma_\mu R q_\beta, \quad O_8 = \frac{3}{2}\bar{D}_\beta\gamma^\mu L b_\alpha \cdot \sum_q e_q\bar{q}_\alpha\gamma_\mu R q_\beta, \\
O_9 &= \frac{3}{2}\bar{D}_\alpha\gamma^\mu L b_\alpha \cdot \sum_q e_q\bar{q}_\beta\gamma_\mu L q_\beta, \quad O_{10} = \frac{3}{2}\bar{D}_\beta\gamma^\mu L b_\alpha \cdot \sum_q e_q\bar{q}_\alpha\gamma_\mu L q_\beta, \\
O_{7\gamma} &= -\frac{e m_b}{8\pi^2}\bar{D}\sigma^{\mu\nu}F_{\mu\nu}(1+\gamma_5)b,
\end{aligned} \tag{2}$$



FIG. 2: Hairpin diagrams for $B \to \phi\rho$ and $B \to \phi\omega$ processes, and similar for $B \to \phi\gamma$.



FIG. 3: The leading diagram from four-quark operators for $\bar{B}^0 \to \phi\gamma$ and $B_s \to \rho\gamma$.

where α and β are the SU(3) color indices and L and R are the left- and right-handed projections operator with $L = (1 - \gamma_5)$, $R = (1 + \gamma_5)$. The sum over q runs over the quark fields that are active at the scale $\mu = \mathcal{O}(m_b)$, i.e., $q = u, d, s, c, b$. The Wilson coefficients evaluated at $\mu = \bar{m}_b(m_b) = 4.4GeV$ from $m_t = 170GeV$ in NDR with $\Lambda^{(5)}_{\bar{M}S} = 225MeV$ ($C_{7\gamma}$ evaluated at $\mu = 5.0$ GeV from $m_t = 170$ GeV with $\alpha_s^{(5)}(M_Z) = 0.118$) are:

$$\begin{aligned} C_1 &= -0.185,\ C_2 = 1.082,\ C_3 = 0.014,\ C_4 = -0.035,\ C_5 = 0.009\ C_6 = -0.041, \\ C_7 &= -0.002/137,\ C_8 = 0.054/137,\ C_9 = -1.292/137,\ C_{10} = 0.263/137, C_{7\gamma} = 0.300. \end{aligned} \tag{3}$$

The radiative decay $\bar{B}^0_d \to \phi\gamma$ has been studied using QCD factorization approach in [12] and using perturbative QCD approach in [13]. There are four diagrams contributing to this decay mode from four-quark operators. The two diagrams with photon emitting from s and $\bar{s}$ cancel with each other. The diagram with photon emitting from the b quark is suppressed by $1/m_b$ compared to the contribution with photon emitting from the light quark in B meson. The dominant contribution



FIG. 4: The electromagnetic dipole operator (denoted by heavy dot) contribution to $\bar{B}^0 \to \phi\gamma$ and $B_s \to \rho\gamma$.

from four-quark operators to $\bar{B}_d^0 \to \phi\gamma$ is illustrated in Fig. 3. In naive factorization, this process are proportional to $C_3 + C_4/3 + C_5 + C_6/3 - 0.5(C_7 + C_8/3 + C_9 + C_{10}/3)$. They have very tiny branching ratios due to the large cancelations between Wilson coefficients: $C_3 \simeq -C_4/3$ and $C_5 \simeq -C_6/3$. This cancellation implies the fact that the dominant diagrams for these decays are the three-gluon hairpin diagrams. To be even worse, the QCD penguins cancel with the electroweak penguins: $C_3+C_4/3+C_5+C_6/3 = -0.002$, $-0.5(C_7+C_8/3+C_9+C_{10}/3) = 0.004$. It is dominated by the photon radiating from the light quark in B meson. Because the ϕ meson is transversely polarized, so the amplitude is suppressed by a power of Λ_{QCD}/m_B further. Factorization properties of the radiative decays $B \to V\gamma$ have been analyzed at the leading power in $1/m_b$ in Ref. [14] (see also [15]) using the soft-collinear effective theory. None of the leading power operators in SCET$_{II}$ will contribute to $\bar{B}_d^0 \to \phi\gamma$, we have to perform the subleading power analysis and this also implies that this mode is suppressed. In [12], the branching ratios of naive factorization and QCD factorization which includes the non-factorizable contributions are:

$$\mathcal{BR}(\bar{B}_d^0 \to \phi\gamma)_{NF} = 3.5 \times 10^{-13}, \tag{4}$$

$$\mathcal{BR}(\bar{B}_d^0 \to \phi\gamma)_{QCDF} = 3.6 \times 10^{-12}. \tag{5}$$

The above results are too rare to be measured in the running B factories BaBar and Belle and may be a probe to detect the new physics. As an example, it can be treated as a probe of R-parity violating couplings (RPV).

In the effective theory, when matching QCD to SCET$_I$ at the leading power, the electromagnetic dipole operators can be factorized into (we use the notation of [14]):

$$\begin{aligned} J^A &= \bar{\mathcal{X}}_{hc}\mathcal{A}^{em}_{\bar{hc}\perp}\Gamma' h_s, \\ J^D &= \bar{\mathcal{X}}_{\bar{hc}}\mathcal{A}^{em}_{hc\perp}\Gamma' h_s. \end{aligned} \tag{6}$$

Including QED in SCET, there is a collinear photon field with unsuppressed interactions with collinear quarks of the same direction. With the additional operator $\mathcal{A}$ overlaps with the transversely polarized vector meson, the J^A and J^D can be matched onto the same operator in SCET$_{II}$:

$$O = \bar{\mathcal{Q}}_s \mathcal{A}^{em}_{c\perp}\mathcal{A}^{em}_{\bar{c}\perp}\Gamma'\mathcal{H}_s. \tag{7}$$

At tree level, these operators correspond to the two diagrams in Fig.4. Using the naive factorization approach the decay amplitude reads:

$$A(\bar{B}_d^0 \to \phi\gamma)_a = \frac{G_F}{\sqrt{2}} V_{tb}^* V_{td} \frac{-2\bar{m}_b e^3 f_\phi Q_d Q_s N_c}{\pi^2 m_\phi m_B \sqrt{2N_c}} C_{7\gamma}\lambda_B^{-1} \times \epsilon^*_{\gamma\alpha} p_{\gamma\beta} p_{\phi\mu} \epsilon^*_{\phi\nu} (g^{\alpha\nu} g^{\beta\mu} - g^{\alpha\mu} g^{\beta\nu} - i\epsilon^{\alpha\beta\mu\nu} \tag{8}$$

TABLE I: Branching ratios of $\bar{B}_d^0 \to \phi\gamma$ (in units of 10^{-11}) and $B_s \to \rho\gamma$ (in units of 10^{-10}) using different $\lambda_{B_{(s)}}^{-1}$

λ_B^{-1}	0.4	0.5	0.6
$\bar{B}_d^0 \to \phi\gamma$(without $O_{7\gamma}$)	0.04	0.06	0.09
$\bar{B}_d^0 \to \phi\gamma$(with $O_{7\gamma}$)	0.86	1.3	1.9
$\lambda_{B_s}^{-1}$	0.4	0.5	0.6
$B_s \to \rho\gamma$ (without $O_{7\gamma}$)	0.45	0.69	1.0
$B_s \to \rho\gamma$(with $O_{7\gamma}$)	5.9	9.2	13

where $\epsilon_\gamma, p_\gamma$ and ϵ_ϕ, p_ϕ are the momentum and polarization for the photon and ϕ meson. $Q_d = Q_s = -1/3, N_c = 3$. λ_B^{-1} is the first inverse moment of the B meson's wave function and defined as

$$\lambda_B^{-1} = \int_0^1 dx \frac{\phi_B(x)}{x}, \tag{9}$$

where x is the momentum fraction carried by the light quark in the B meson. The neutral vector meson is generated by the intermediate photon, so we can view the two diagrams in Fig. 4 as $B_{d,s} \to \gamma\gamma$ [16] at first. The second diagram can be related to the first one by the crossing symmetry, so it gives the same contribution as the first one.

There are many candidates for B meson distribution amplitudes [5, 17], but here only λ_B^{-1} is involved and we use three values for this parameter: $\lambda_B^{-1} = 0.4, 0.5, 0.6$. We adopt other parameters the same as [12, 13], then numerical results for the branching ratios of these two modes are given in Table I. For $B_s \to \rho\gamma$, there is also a contribution from the tree operators O_1 and O_2. But this contribution is suppressed by the CKM matrix elements and roughly is of order 10^{-12}, so we neglect it in the Table. We find that the branching ratios are enhanced sizably by the new contribution relative to the result from QCD factorization approach. Although it is hardly to be measured at BaBar and Belle, we state that the contribution of order 10^{-11} for $\bar{B}_d^0 \to \phi\gamma$ would not be the evidence of the activity of new physics. As for $B_s \to \rho\gamma$, $\mathcal{BR} \sim (6-16) \times 10^{-10}$ which might be detected at the LHCb experiment.

For $B \to \rho\phi$ and $B \to \omega\phi$, they are very similar to the above radiative decays. These processes can also only be produced via the so-called hair-pin diagrams as in Fig. 2 [10] which are highly suppressed and the branching ratios in generalized factorization [2] are:

$$\begin{aligned} \mathcal{BR}(B^+ \to \rho^+\phi) &= 0.04 \times 10^{-7}, \\ \mathcal{BR}(B^0 \to \rho^0\phi) &= 0.02 \times 10^{-7}, \\ \mathcal{BR}(B^0 \to \omega\phi) &= 0.02 \times 10^{-7}. \end{aligned} \tag{10}$$

In this calculations, the electromagnetic dipole operator $O_{7\gamma}$ is also neglected. This operator contributes only to the transversely polarized amplitude. The contribution can be factorized into [9]:

$$\mathcal{A}(B \to V_1 V_2) = i m_{V_2} m_B 2 T_1^{V_1}(0) f_{V_2} a_{V_2} \times \Big(- \frac{2\alpha_{em}}{3\pi} \Big) C_{7\gamma} \frac{m_B m_b}{m_{V_2}^2}, \tag{11}$$

where V_2 is a neutral vector meson produced by the photon. a_{V_2} is a constant which depends on the quark-flavor composition of V_2: $a_\phi = -1/2$, $a_\rho = 3/(2\sqrt{2})$. The factor $\sqrt{2}$ arises from the normalization of the ρ^0 meson. $T_1^{V_1}(0)$ is the QCD tensor form factor of $B \to V_1$, f_{V_2} and m_{V_2} is the decay constant and mass of V_2. The amplitude takes the given value only when both V_1 and V_2 have negative helicity, but is zero otherwise.

The four-quark operators in Fig.1(b) from the effective Hamiltonian could give corrections to the above expression and they have been computed in next-to-leading order in the context of factorization of exclusive radiative B decays [18, 19]. After incorporating this contribution, the Wilson coefficient $C_{7\gamma}$ is replaced by $\mathcal{C}_7'$ [18]. The tensor form factor $T_1^{V_1}(0)$ is a nonperturbative parameter. In PQCD approach, it can be factorized into the convolution of the wave function and the hard part. We take the results from the PQCD calculations [20]:

$$T_1^{B\to\rho}(0) = 0.41, \;\; T_1^{B\to\omega}(0) = 0.38. \tag{12}$$

Using the above inputs and $f_\phi = 0.254\text{GeV}, m_\phi = 1.02\text{GeV}, m_B = 5.28\text{GeV}, |V_{td}| = 0.008, |V_{tb}| = 1.0$, we get

$$\begin{aligned} \mathcal{BR}(B^+ \to \rho^+\phi) &= 3.0^{+0.5}_{-0.5}(1.6) \times 10^{-9}, \\ \mathcal{BR}(B^0 \to \rho^0\phi) &= 1.4^{+0.2}_{-0.2}(0.77) \times 10^{-9}, \\ \mathcal{BR}(B^0 \to \omega\phi) &= 1.2^{+0.2}_{-0.2}(0.66) \times 10^{-9}. \end{aligned} \tag{13}$$

Results in the parentheses are performed by using the leading order Wilson coefficient for $C_{7\gamma}$, while the others are evaluated by using $|\mathcal{C}_7'|^2 = 0.165^{+0.018}_{-0.017}$ [18]. From the results, we can see that the new contribution is comparable to the original one which may imply that the large transverse polarization in this mode. Compared with the experimental result by BaBar collaboration [21]:

$$\mathcal{BR}(B^0 \to \phi\omega) = 0.1 \pm 0.5 \pm 0.1 (< 1.2) \times 10^{-6}, \tag{14}$$

we can see that these modes can hardly be measured in the running B factories. If we replace B by B_s ($V_{td} \to V_{ts}$) and assume SU(3) symmetry to give an estimate, the corresponding processes

$B_s \to \phi\rho^0$ and $B_s \to \phi\omega$ can be enhanced to 10^{-7} which may be measured at the future LHCb experiment.

In summary, we have studied the enhancement of $\bar{B}_d^0 \to \phi\gamma$ and $B_s \to \rho\gamma$ due to the electromagnetic dipole operators. This effect could produce sizable contributions to $\bar{B}_d^0 \to \phi\gamma$ to make a branching ratio of about 1.0×10^{-11}, which is larger than the previous study using QCD factorization approach. Although it is hardly to be measured at BaBar and Belle, we conclude that the contribution of order 10^{-11} would not be the evidence of the activity of new physics. As for $B_s \to \rho\gamma$, the branching ratio is $(6 - 16) \times 10^{-10}$. For $B \to \rho\phi$ and $B \to \omega\phi$, the new effect also give a large contribution, of order of 10^{-9}, to the transverse helicity amplitude which is comparable to the longitudinal amplitude. So after incorporating this effect, the polarization can be changed sizably from the naive power counting for $B \to VV$. These effects may be detected in the future LHCb experiment.

---

[1] M. Wirbel, B. Stech, and M. Bauer, Z. Phys. C **29**, 637 (1985); M. Bauer, B. Stech, and M. Wirbel, Z. Phys. C **34**, 103 (1987); L.-L. Chau, H.-Y. Cheng, W.K. Sze, H. Yao, and B. Tseng, Phys. Rev. D **43**, 2176, (1991), Erratum: **58**, 019902 (1998).

[2] A. Ali, G. Kramer and C.-D. Lü, Phys. Rev. D **58**, 094009 (1998); C.-D. Lü, Nucl. Phys. B. Proc. Supp. **74**, 227 (1999).

[3] Y.-H. Chen, H.-Y. Cheng, B. Tseng, and K.-C. Yang, Phys. Rev. D **60**, 094014 (1999); H.-Y. Cheng and K.-C. Yang, Phys. Rev. D **62**, 054029 (2000).

[4] M. Beneke, G. Buchalla, M. Neubert, and C.T.Sachrajda, Phys. Rev. Lett. **83**, 1914 (1999); Nucl. Phys.**B 591**, 313 (2000); Nucl. Phys. **B 606**, 245 (2001).

[5] Y.Y. Keum, H.-n. Li, and A.I. Sanda, Phys. Lett. B **504**, 6 (2001); Phys. Rev. D **63**, 054008 (2001); Y.Y. Keum and H.-n. Li, Phys. Rev. D **63**, 074006 (2001); C.-D. Lu, K. Ukai and M.-Z. Yang, Phys. Rev. D **63**, 074009 (2001); C.-D. Lu and M.-Z. Yang, Eur. Phys. J. C **23**, 275 (2002).

[6] C.W. Bauer, S. Fleming and M.E. Luke, Phys. Rev. D **63**, 014006 (2001); C.W. Bauer, S. Fleming, D. Pirjol and I.W. Stewart, Phys. Rev. D **63**, 114020 (2001); C.W. Bauer and I.W. Stewart, Phys. Lett. B **516**, 134 (2001); J. Chay and C. Kim, Phys. Rev. D **65**, 114016 (2002); M. Beneke, A.P. Chapovsky, M. Diehl and T.Feldmann, Nucl. Phys. **B 643**, 431 (2002); C.W. Bauer, D. Pirjol, I.Z. Rothstein and I.W. Stewart, Phys. Rev. D 70, 054015 (2004).

[7] A.L. Kagan, Phys. Lett. B **601**,151 (2004); H.-n. Li and S. Mishima, Phys. Rev. D **71**, 054025 (2005); H.-n. Li, Phys. Lett. B **622**, 63 (2005).

[8] E. Barberio *et al.* [The Heavy Flavor Averaging Group], arXiv: hep-ex/0603003.

[9] M. Beneke, J. Rohrer and D. Yang, Phys. Rev. Lett. **96**, 141801 (2006)

[10] D.S. Du and Z.Z. Xing, Phys. Lett. B **312**, 199 (1993).

[11] For a review, see G. Buchalla, A.J. Buras and M.E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1006).

[12] Y.D. Yang, Eur. Phys. J. C **36**, 97 (2004).

[13] Y. Li and C.D. Lu, arXiv: hep-ph/0605220.

[14] T. Becher, R.J. Hill and M. Neubert, Phys. Rev. D **72**, 094017 (2005).

[15] J.G. Chay and C. Kim, Phys. Rev. D **68**, 034013 (2003).

[16] S.W. Bosch and G. Buchalla, JHEP, **0208**, 054, (2002)

[17] A.G. Grozin and M. Neubert, Phys. Rev. D **55**, 272 (1997); H. Kawamura, J. Kodaira, C.F. Qiao and K. Tanaka, Phys. Lett. B **523**, 111 (2001); Erratum-ibid. **536**, 344 (2002); Mod. Phys. Lett. A **18**, 799 (2003); see also T. Kurimoto, arXiv: hep-ph/0605112.

[18] M. Beneke, Th. Feldmann and D. Seidel, Nucl. Phys. **B 612**, 25 (2001).

[19] S.W. Bosch and G. Buchalla, Nucl. Phys. **B 621**, 459 (2002).

[20] C.-D. Lu and M.-Z. Yang, Eur. Phys. J. C **28**, 515 (2003).

[21] The BABAR Collaboration, B. Aubert, *et al.* arXiv: hep-ex/0605017.